


For use with a Puffer and Spacer

Name: \_\_\_\_\_

Date of birth: \_\_\_\_\_



Photo

Child can self administer medication if well enough.

Child needs to pre-medicate prior to exercise

Confirmed triggers:

Family/emergency contact name(s):

Work Ph: \_\_\_\_\_

Home Ph: \_\_\_\_\_

Mobile Ph: \_\_\_\_\_

Plan prepared by Dr or NP:

I hereby authorise medications specified on this plan to be administered according to the plan.

Signed:

Date: \_\_\_\_\_

Date of next review: \_\_\_\_\_



- Assemble Spacer
- Remove cap from puffer
- Shake puffer well
- Attach puffer to end of spacer
- Place mouthpiece of spacer in mouth and ensure lips seal around it
- Breath out gently into the spacer
- Press down on puffer canister once to fire medication into spacer
- Breathe in and out normally for 4 breaths (keeping your mouth on the spacer)

## MILD TO MODERATE SIGNS

- Minor difficulty breathing
- May have a cough
- May have a wheeze

## ACTION FOR MILD TO MODERATE ASTHMA FLARE UP

- 1 Sit the person upright.
  - Stay with person and be calm and reassuring
- 2 Give..... separate puffs of Airomir, Asmol or Ventolin
  - Shake puffer before each puff
  - Put 1 puff into the spacer at a time
  - Take 4 breaths from the spacer between each puff
- 3 Wait 4 minutes.
  - If there is no improvement, repeat step 2

**If there is still no improvement follow the Asthma First Aid Plan for severe / life-threatening asthma attacks below**

Mild to moderate symptoms do not always present before severe or life-threatening symptoms

## SEVERE SIGNS

- Cannot speak a full sentence
- Sitting hunched forward
- Tugging in of skin over chest or throat
- May have a cough or wheeze
- Obvious difficulty breathing
- Lethargic
- Sore tummy (young children)

## LIFE-THREATENING SIGNS

- Unable to speak or 1-2 words
- Collapsed / Exhausted
- Gasping for breath
- May no longer have a cough or wheeze
- Drowsy/ Confused / Unconscious
- Skin discolouration (blue lips)

## ACTION FOR SEVERE / LIFE-THREATENING ASTHMA ATTACK

- 1 Sit the person upright. Be calm and reassuring. Do not leave them alone.
- 2 Phone ambulance: Triple Zero (000).
- 3 Give ..... separate puffs of Airomir, Asmol or Ventolin
  - Shake puffer before each puff
  - Put 1 puff into the spacer at a time
  - Take 4 breaths from the spacer between each puff
- 4 Wait 4 minutes.
- 5 Keep giving ..... puffs every 4 minutes until emergency assistance arrives.

Commence CPR at any time if person is unresponsive and not breathing normally. Blue reliever medication is unlikely to harm, even if the person does not have asthma.

## IF UNCERTAIN WHETHER IT IS ANAPHYLAXIS OR ASTHMA

- Give adrenaline autoinjector FIRST, then asthma reliever.
- If someone with known food or insect allergy suddenly develops severe asthma like symptoms, give adrenaline autoinjector FIRST, then asthma reliever.

Anaphylaxis: Y  N  Type of autoinjector: \_\_\_\_\_