

Patient name: _____ Date: DD / MM / YYYY
Plan prepared by: _____ Signed: _____

ALLERGEN MINIMISATION

- ☐ Minimising exposure to confirmed allergen/s may assist to reduce symptoms in some people.
For information go to www.allergy.org.au/patients/allergy-treatments/allergen-minimisation

THUNDERSTORM ASTHMA

- ☐ Try to stay indoors just before, during and just after thunderstorms in pollen seasons if allergic to pollen.
Use preventer treatments such as intranasal corticosteroid sprays or combined intranasal corticosteroid/antihistamine sprays. Consider allergen immunotherapy (see below). If you also have asthma, use asthma preventers regularly.
For information go to www.allergy.org.au/patients/asthma-and-allergy/thunderstorm-asthma


MEDICATIONS

- ☐ Intranasal corticosteroid spray: _____
☐ 1 or ☐ 2 times/day/nostril for _____ weeks or _____ months or ☐ continuous
☐ Additional instructions: _____
or
☐ Combined intranasal corticosteroid/antihistamine spray:
☐ 1 or ☐ 2 times/day/nostril for _____ weeks or _____ months or ☐ continuous
☐ Additional instructions: _____

Note:

- It is important to use these sprays correctly. See instructions below and directions for use.
- Onset of benefit may take days, so these sprays must be used regularly and should not be stopped every few weeks.
- If significant pain or bleeding occurs contact your doctor.
- Some treatments mentioned above require a prescription.

1. Prime the spray device according to manufacturer's instructions (for the first time or after a period of non-use).
2. Shake the bottle before each use.
3. Blow nose before spraying if blocked by mucus.
4. Tilt head slightly forward and gently insert nozzle into nostril.
5. Aim the nozzle away from the middle of the nose (septum) and direct nozzle into the nasal passage (not towards tip of nose, but in line with the roof of the mouth).
6. Avoid sniffing hard during or after spraying.



- ☐ Oral non-sedating antihistamine tablet: _____ Dose _____ mL/mg ☐ 1 or ☐ 2 times/day
☐ Additional instructions: _____
- ☐ Intranasal antihistamine sprays: _____ ☐ 1 or ☐ 2 times/day
☐ Additional instructions: _____
- ☐ Saline nasal ☐ spray or ☐ irrigation _____ ☐ _____ times/day or ☐ as needed
☐ Use 10 minutes prior if used with intranasal corticosteroid spray
- ☐ Decongestant: _____ ☐ nasal spray _____ times/day or ☐ tablet
Dose _____ tablets _____ times/day for up to three days (not more than one course/month)
- ☐ Eye drops or ointments: _____
- ☐ Other medications: _____

For information and links to animation videos go to www.allergy.org.au/patients/allergic-rhinitis-hay-fever-and-sinusitis

ALLERGEN IMMUNOTHERAPY

If allergen immunotherapy has been initiated by a clinical immunology/allergy specialist or other doctor, it is important to follow the treatment as prescribed. Contact your doctor if you have any questions or concerns.
For information go to www.allergy.org.au/patients/allergy-treatments/immunotherapy