



# Consent form for COVID-19 vaccination

Before you fill out this form, make sure you read the information sheet on the vaccine you will be getting: Vaxzevria (AstraZeneca), Comirnaty (Pfizer), Spikevax (Moderna), or Nuvaxovid (Novavax).

Last updated: 2 February 2022

#### **About COVID-19 vaccination**

People who have a COVID-19 vaccination course have a much lower chance of getting sick from COVID-19.

There are four brands of vaccine in use in Australia. All are effective and safe.

#### You can have:

- AstraZeneca or Novavax if you are 18 years or over
- Moderna if you are 12 years or over
- Pfizer if you are 5 years or over\*

Pfizer, Moderna, or Novavax are preferred over AstraZeneca for adults under 60 years of age.

Most people require two doses initially. This is called the primary course.

People with severe immunocompromise may require a third primary dose to bring their immune response up to optimal levels.

People aged 16 years or over, including people with severe immunocompromise and pregnant women, should have a booster dose three months or more after their primary course, to prolong their protection against COVID-19.

- People aged 16 and over can have a booster dose of Pfizer.
- People aged 18 and over can have Pfizer or Moderna as a booster.
- AstraZeneca is not preferred but can be used in some instances in people over 18 years.

For more information visit the Department of Health COVID-19 vaccine website: <a href="https://www.health.gov.au/covid-19-vaccines">www.health.gov.au/covid-19-vaccines</a>

Name:						
Medicare number:						

<sup>\*</sup>There is a separate consent form available for children aged 5 to 11 years.

Medical experts have studied COVID-19 vaccines to make sure they are safe. Most side effects are mild. They may start on the day of vaccination and last for one or two days. As with any vaccine or medicine, there may be rare or unknown side effects.

A very rare side effect after AstraZeneca is called thrombosis with thrombocytopenia syndrome, or TTS. This means blood clotting (thrombosis) with low blood platelet levels (thrombocytopenia). TTS does not happen after Pfizer or Moderna.

Myocarditis and pericarditis (heart inflammation) have been reported following Pfizer and Moderna. Most cases have been mild and people have recovered quickly.

Tell your health care provider if you have any side effects after vaccination that you are worried about.

You may be contacted by SMS or email in the week after you have the vaccine to see how you are feeling.

Some people may get COVID-19 after vaccination. You must still follow all public health advice in your state or territory to stop the spread of COVID-19, including:

- keep your distance stay at least 1.5 metres away from other people
- wash your hands often with soap and water, or use hand sanitiser
- wear a mask
- stay home if you are unwell with cold or flu-like symptoms and arrange to get a COVID-19 test.

By law, the person giving your vaccination must record it on the Australian Immunisation Register. You can view your vaccination record online through your:

- Medicare account
- MyGov account
- My Health Record account (you can register for this with a Medicare number or an Individual Healthcare Identifier).

## How your information is used

For information on how your personal details are collected, stored and used, visit www.health.gov.au/using-our-websites/privacy/privacy-notice-for-covid-19-vaccinations.

If you are getting your vaccination in a pharmacy, the pharmacy must share some of your personal information with the Pharmacy Programs Administrator. This is so the pharmacy can claim payment from the Australian Government. More information about why this is needed and what information is shared is provided at the link above.

### On the day you have your vaccine

Before you get vaccinated, tell the person giving you the vaccination if you:

- have had an allergic reaction, particularly a severe allergic reaction (anaphylaxis), to:
  - o a previous dose of a COVID-19 vaccine
  - o an ingredient of a COVID-19 vaccine
  - o other vaccines or medications
- are immunocompromised. This means that you have a weakened immune system that
  makes it harder for you to fight diseases. You can still have a COVID-19 vaccine but talk to
  your doctor about when is the best time to get your vaccine. This will depend on your
  condition and your treatment.

Name:						
Medicare number:						

# **Consent Checklist**

Yes	No	
		Have you had an allergic reaction to a previous dose of a COVID-19 vaccine?
		Have you had anaphylaxis to another vaccine or medication?
		Have you had a serious adverse event, that following expert review by an experienced immunisation provider or medical specialist was attributed to a previous dose of a COVID-19 vaccine (and did not have another cause identified)?
		Have you ever had mastocytosis (a mast cell disorder) which has caused recurrent anaphylaxis?
		Have you had COVID-19 before?
		Do you have a bleeding disorder?
		Do you take any medicine to thin your blood (an anticoagulant therapy)?
		Do you have a weakened immune system (immunocompromised)?
		Are you pregnant? #
		Have you been sick with a cough, sore throat, fever or are feeling sick in another way?
		Have you had a COVID-19 vaccination before?
		Have you received any other vaccination in the last 7 days?
Relev	ant or	nly for those receiving AstraZeneca:
		Have you ever been diagnosed with capillary leak syndrome?
		Have had thrombosis (clotting) together with thrombocytopenia (low platelets) within 42 days after having a previous dose of AstraZeneca?
		Have you ever had cerebral venous sinus thrombosis? *
		Have you ever had heparin-induced thrombocytopenia? *
		Have you ever had blood clots in the abdominal veins (splanchnic veins)? *
		Have you ever had antiphospholipid syndrome associated with blood clots? *
		Are you under 60 years of age? *
Novava	x or As	oderna are the preferred vaccines for pregnant women. If these vaccines are not available, straZeneca can be considered. More information see: <a href="https://www.health.gov.au/initiatives-/covid-19-vaccines/who-can-get-vaccinated/pregnant-women">https://www.health.gov.au/initiatives-/covid-19-vaccines/who-can-get-vaccinated/pregnant-women</a>
		erna, or Novavax are the preferred vaccines for people in these groups. If these vaccines ble, AstraZeneca can be considered if the benefits of vaccination outweigh the risks.
For mor		rmation, see <a href="https://www.health.gov.au/initiatives-and-programs/covid-19-vaccines/advice-ts">https://www.health.gov.au/initiatives-and-programs/covid-19-vaccines/advice-ts</a>

Name:

Medicare number:

Releva	ant only	y for those receiving	Pfize	r or N	1odei	ma:								
		Have you been diag	-		th my	ocard	itis aı	nd/or	peric	arditi	s afte	er a p	orevio	us
		Have you had myo	cardit	is or	peric	arditis	withi	n the	past	three	mor	ths?		
		Do you currently ha	ive a	cute r	heun	natic f	ever	or acı	ute rh	euma	atic h	eart	disea	se?
		Do you have severe	e hea	ırt fail	ure?									
howeve of vacci	er you s ination	ed <b>Yes</b> to any of the ab should talk to your GP, and whether any addit	immu	ınisati	on sp	ecialis	or ca	rdiolo						
·		2 February 2022												
Pati	ent	information												
Name	e:													
Medic	care nu	ımber:												
	dual H licable	ealth Identifier (IHI) :												
Date	of birth	1:												
Addre	ess:													
Phone	e conta	act number:												
Email	addre	ess:												
Gend	er:													
Langı	uage s	poken at home:												
Coun	try of b	oirth:												
Are vo	u Abor	iginal and/or Torres	Strait	Islan	ider?									
☐ Yes ☐ Yes ☐ Yes ☐ No	s, Abor s, Torre s Abori	iginal only es Strait Islander onl ginal and Torres Stra to answer	y											
Next	of kin (	in case of emergeno	;y):											
Name	e:													
Phone	e conta	act number:												

Name:						
Medicare number:						

Con	sent to receive	COVID-	19 vaccine							
	I confirm I have revaccination.	eceived a	and understood information provided to me on COVID-19							
		ny other s	of the above conditions apply to me, or I have discussed these special circumstances with my regular health care provider er.							
☐ I agree to receive a course of COVID-19 vaccine / I agree to receive a booster of COVID-19 vaccine										
Pati	ent's name:									
Pati	ent's signature:									
Dat	e:									
	-	•	guardian or substitute decision-maker, and agree to the patient named above.							
	ent/guardian/subst ision-maker's nam									
Parent/guardian/substitute decision maker's signature:										
Dat	e:									

Name:						
Medicare number:						

# For provider use:

#### Dose 1:

Date vaccine administered:	
Time received:	
COVID-19 vaccine brand administered:	
Batch no:	
Serial no:	
Site of vaccine injection:	
Name of vaccination service provider:	
Dose 2:	
Date vaccine administered:	
Time received:	
COVID-19 vaccine brand administered:	
Batch no:	
Serial no:	
Site of vaccine injection:	
Name of vaccination service provider:	
Dose 3 or booster*:	
Date vaccine administered:	
Time received:	
COVID-19 vaccine brand administered:	
Batch no:	
Serial no:	
Site of vaccine injection:	
Name of vaccination service provider:	
	primary dose of COVID-19 vaccine in individuals
	recommendations on the use of a booster dose
of COVID-19 vaccine, and ATAGI recommenda	ations on the use of Spikevax Moderna as a

booster dose.

Last updated: 2 February 2022

Name:						
Medicare number:						