




**For use with a Puffer and Spacer**

Name: \_\_\_\_\_

Date of birth: \_\_\_\_\_



Photo

Child can self administer medication if well enough.  
Child needs to pre-medicate prior to exercise

Confirmed triggers:

Family/emergency contact name(s):

Work Ph: \_\_\_\_\_

Home Ph: \_\_\_\_\_

Mobile Ph: \_\_\_\_\_


Plan prepared by Dr or NP:

**I hereby authorise medications specified on this plan to be administered according to the plan.**

Signed:

Date: \_\_\_\_\_

Date of next review: \_\_\_\_\_



- Assemble Spacer
- Remove cap from puffer
- Shake puffer well
- Attach puffer to end of spacer
- Place mouthpiece of spacer in mouth and ensure lips seal around it
- Breath out gently into the spacer
- Press down on puffer canister once to fire medication into spacer
- Breathe in and out normally for 4 breaths (keeping your mouth on the spacer)

## MILD TO MODERATE SIGNS

- Minor difficulty breathing
- May have a cough
- May have a wheeze

## ACTION FOR MILD TO MODERATE ASTHMA FLARE UP

- 1** Sit the person upright.
  - Stay with person and be calm and reassuring
- 2** Give 4 separate puffs of Airomir, Asmol or Ventolin
  - Shake puffer before each puff
  - Put 1 puff into the spacer at a time
  - Take 4 breaths from the spacer between each puff
- 3** Wait 4 minutes.
  - If there is no improvement, repeat step 2

**If there is still no improvement follow the Asthma First Aid Plan for severe / life-threatening asthma attacks below**

**Mild to moderate symptoms do not always present before severe or life-threatening symptoms**

## SEVERE SIGNS

- Cannot speak a full sentence
- Sitting hunched forward
- Tugging in of skin over chest or throat
- May have a cough or wheeze
- Obvious difficulty breathing
- Lethargic
- Sore tummy (young children)

## LIFE-THREATENING SIGNS

- Unable to speak or 1-2 words
- Collapsed / Exhausted
- Gasping for breath
- May no longer have a cough or wheeze
- Drowsy/ Confused / Unconscious
- Skin discolouration (blue lips)

## ACTION FOR SEVERE / LIFE-THREATENING ASTHMA ATTACK

- 1 Sit the person upright. Be calm and reassuring. Do not leave them alone.**
- 2 Phone ambulance: Triple Zero (000).**
- 3 Give 4 separate puffs of Airomir, Asmol or Ventolin**
  - Shake puffer before each puff
  - Put 1 puff into the spacer at a time
  - Take 4 breaths from the spacer between each puff
- 4 Wait 4 minutes.**
- 5 Keep giving 4 puffs every 4 minutes until emergency assistance arrives.**

**Commence CPR at any time if person is unresponsive and not breathing normally.**  
Blue reliever medication is unlikely to harm, even if the person does not have asthma.

## IF UNCERTAIN WHETHER IT IS ANAPHYLAXIS OR ASTHMA

- Give adrenaline autoinjector **FIRST**, then asthma reliever.
- If someone with known food or insect allergy suddenly develops severe asthma like symptoms, give adrenaline autoinjector **FIRST**, then asthma reliever.

Anaphylaxis: Y  N  Type of autoinjector: \_\_\_\_\_