


Name: \_\_\_\_\_  
 Date of birth: \_\_\_\_\_



Confirmed allergens:

Family/emergency contact name(s):  
 \_\_\_\_\_  
 \_\_\_\_\_

Work Ph: \_\_\_\_\_  
 Home Ph: \_\_\_\_\_  
 Mobile Ph: \_\_\_\_\_

Plan prepared by medical or nurse practitioner:

I hereby authorise medications specified on this plan to be administered according to the plan

Signed: \_\_\_\_\_  
 Date: \_\_\_\_\_  
 Action Plan due for review: \_\_\_\_\_

Note: This ASCIA Action Plan for Allergic Reactions is for people with mild to moderate allergies, who need to avoid certain allergens

For people with severe allergies (and at risk of anaphylaxis) there are ASCIA Action Plans for Anaphylaxis, which include adrenaline (epinephrine) autoinjector instructions

Instructions are also on the device label

Note: All EpiPen®s should be held in place for 3 seconds regardless of instructions on device label

## SIGNS OF MILD TO MODERATE ALLERGIC REACTION

- Swelling of lips, face, eyes
- Hives or welts
- Tingling mouth
- Abdominal pain, vomiting (these are signs of anaphylaxis for insect allergy)

## ACTION FOR MILD TO MODERATE ALLERGIC REACTION

- For insect allergy - flick out sting if visible
- For tick allergy - freeze dry tick and allow to drop off
- Stay with person and call for help
- Give other medications (if prescribed).....
- Phone family/emergency contact

**Mild to moderate allergic reactions (such as hives or swelling) may not always occur before anaphylaxis**

## WATCH FOR ANY ONE OF THE FOLLOWING SIGNS OF ANAPHYLAXIS (SEVERE ALLERGIC REACTION)

- **Difficult/noisy breathing**
- **Swelling of tongue**
- **Swelling/tightness in throat**
- **Wheeze or persistent cough**
- **Difficulty talking and/or hoarse voice**
- **Persistent dizziness or collapse**
- **Pale and floppy (young children)**

## ACTION FOR ANAPHYLAXIS

**1 Lay person flat - do NOT allow them to stand or walk**

- If unconscious, place in recovery position

- If breathing is difficult allow them to sit



**2 Give adrenaline (epinephrine) autoinjector if available**

**3 Phone ambulance - 000 (AU) or 111 (NZ)**

**4 Phone family/emergency contact**

**5 Transfer person to hospital for at least 4 hours of observation**

**If in doubt give adrenaline autoinjector**

**Commence CPR at any time if person is unresponsive and not breathing normally**

**ALWAYS give adrenaline autoinjector FIRST if available, and then asthma reliever puffer** if someone with known asthma and allergy to food, insects or medication has **SUDDEN BREATHING DIFFICULTY** (including wheeze, persistent cough or hoarse voice) even if there are no skin symptoms

Asthma reliever medication prescribed:  Y  N